

FY18 USWBSI Project Abstract

PI: Paul Esker

PI's E-mail: pde6@psu

Project ID: FY18-IM-008

ARS Agreement #: 59-0206-8-210

Research Category: MGMT

Duration of Award: 1 Year

Project Title: Integrated Management of Fusarium Head Blight in Wheat in Pennsylvania.

PROJECT 1 ABSTRACT

(1 Page Limit)

In Pennsylvania, winter wheat is produced on approximately 150,000 to 200,000 acres, with yields of 65 bu/ac, on average, providing \$40,000,000 to \$70,000,000 to the local economy annually. Nonetheless, questions remain regarding the efficacy of foliar fungicides in an integrated program, as many producers in PA apply fungicides based on growth stage without necessarily quantifying the agronomic, biological or economic efficacy of those applications. New research, as part of the Integrated Management (IM)-Coordinated Project and the Uniform Fungicide Trial (UFT) Project, will help provide important local data to quantify efficacy and improve management recommendations for control of Fusarium head blight (FHB), as well as contributing to national efforts on both best management practices and disease forecasting. IM and UFT trials will be conducted at the Russell E. Larson Agricultural Research Center at Rock Springs and the Southeast Agricultural Research & Extension Center at Landisville, PA, respectfully. A split plot experimental design will be used for the IM trial, while a randomized completed block design will be used for the UFT. Plot size for both trials will be 7 rows by 18 feet, with a harvested length of 14 feet. For the IM trial, the whole plot factor will be winter wheat variety (susceptible and resistant), and the subplot will be the foliar fungicide treatment. For the UFT the treatments are different foliar fungicide products and application timing. A susceptible winter wheat variety will be used. Plots will be established and maintained following local recommendations. FHB incidence and severity will be rated on 60-100 spikes per plot at the soft dough growth stage (Feekes 11.2), as well as the presence and flag leaf severity (as a percentage) of foliar diseases. Two subsamples of grain from each plot will be obtained, with one used to determine the percentage of Fusarium damaged kernels and the other that will be submitted for DON analysis at one of the USWBSI-fungicide laboratories. factor, the PI will where applicable). Data generated from these studies will be used to determine optimal management techniques both locally for the growers of Pennsylvania, as well as part of the larger national efforts to establish effective, economical approaches to FHB and DON management and to improve validation of current forecasting models for FHB. Results will be disseminated through field days, grower meetings, workshops, newsletters, fact sheets and become part of the PSU Agronomy Guide.