



U.S. Wheat and Barley Scab Initiative

FHB Tool Talk

Dear Extension Specialists, Crop Consultants, and Grower Organizations,

It's time to start making plans for fungicide applications to manage Fusarium head blight (FHB) and deoxynivalenol (DON) in winter wheat and barley.

Step 1: Time it right for maximum control. Use the USWBSI's fungicide timing postcards to identify the best time for making an application.

In winter wheat, the best time to apply a fungicide for FHB and DON management is at Feekes growth stage 10.51, which is when **50% of the main tillers have reached early anthesis (flowering) and up to seven days later**. In addition, USWBSI-supported research has shown that fungicide applications made up to 7 days after Feekes growth stage 10.51 may provide similar management of FHB and DON. Be sure to read and follow the fungicide product label to ensure that applications after Feekes growth stage 10.51 are permitted.

FHB Fungicide Timing: Wheat
(Hard & Soft Winter, Spring)


Feekes 10.51 (Early Anthesis)
The optimum time to apply a single fungicide to suppress Fusarium head blight (scab) and reduce deoxynivalenol (DON) is at Feekes growth stage 10.51 (50% of the main tillers have reached early anthesis/flowering) and up to seven days later.

emerged anthers

For more information, go to scabusa.org or contact your local Extension office.

For winter barley, the best time for a fungicide application for FHB and DON management is about **6 days after** full heading (Feekes growth stage 10.5). Application before heads are fully emerged will result in poor suppression of FHB and DON. Be sure to read and follow the

fungicide product label to ensure that applications after Feekes growth stage 10.5 are permitted.



Step 2: **Determine if an application is warranted.** Consult the [FHB Risk Tool](#) to determine if the risk is high in your region.

- The FHB Risk Tool is now active for the 2026 growing season!

Step 3: **Choose a fungicide with care.** Not all fungicides registered on small grain crops are effective in managing FHB. See the [Fungicide Efficacy Guide for Control of Wheat Diseases](#) on the Crop Protection Network to help pick the best fungicide options for managing FHB.

- See figure below** for an efficacy comparison among fungicides obtained from multi-state research funded by the USWBSI. The results include data from over 20 trials conducted in 24 states over two years. Data were compiled, analyzed, and summarized by Dr. Pierce Paul’s lab, The Ohio State University.

i. Fungicide rates.

| | |
|--------------------------------------|-----------------------------|
| Prosaro | 6.5 fl oz/A |
| Caramba | 13.5 fl oz/A |
| Miravis Ace | 13.7 fl oz/A |
| Prosaro Pro | 10.3 fl oz/A |
| Sphaerex | 7.3 fl oz/A |
| Miravis Ace followed by Prosaro Pro | 13.7 fl oz/A 10.3 fl oz/A |
| Miravis Ace followed by Sphaerex | 13.7 fl oz/A 7.3 fl oz/A |
| Miravis Ace followed by Tebuconazole | 13.7 fl oz/A 4.0 fl oz/A |




Fig.1: Reduction of DON (ppm) as a result of fungicide treatments relative to the untreated check.

Step 4: Make application.

- **Configure your ground sprayer to optimize spray coverage.** This includes using a spray volume of 10-20 gallons per acre, utilizing fine to medium size droplets, positioning nozzles 8-10 inches above the heads, and positioning nozzles forward and backward at 30 to 45 degrees down from horizontal. See [“Optimizing Fungicide Use for Fusarium Head Blight \(Scab\) and Associated Mycotoxins”](#) on the Crop Protection Network and [“Ground Application of Fungicides for Fusarium Head Blight Management”](#) by North Dakota State University for more information.
- **Applying a fungicide twice for FHB management?** Results from field research trials funded by the USWBSI have shown that additional reduction in FHB and DON can occur with two treatment programs. More research is needed to better understand the circumstances in which this practice would be economical. Also, it is important to be mindful of product label restrictions that might not allow a two-treatment application in some instances.

Thank you for reading!

If this email was forwarded to you and you would like to receive FHB Tool Talk e-newsletters, [subscribe today](#).

*In order to target a wide range of grain classes and production regions, you may have received this message sooner than you needed. Please bookmark this email and refer back to it when the timing is correct for your region/grain class.



More FHB Resources for [Extension, Crop Consultants, and Grower Organizations](#).

Copyright © 2026 U.S. Wheat and Barley Scab Initiative. All rights reserved.

*U.S. Wheat and Barley Scab Initiative
495 Borlaug Hall
1991 Upper Buford Circle
St. Paul, MN 55108
nfo@scabusa.org*

Want to change how you receive these emails?

[Update subscription preferences.](#)

To unsubscribe from this list, email fbtooltalk_unsubscribe@scabusa.org with the word unsubscribe.

