

PI: Bruce Bleakley**Project ID: 0405-BL-063****Research Area: CBC****Project Title: Management of Fusarium Head Blight with Biological Control Agents.****PI's E-mail: Bruce_Bleakley@sdstate.edu****FY03 ARS Agreement #: 59-0790-0-061****Duration of Award: 1 Year**

PROJECT 1 ABSTRACT
(1 Page Limit)

A field-screening trial for the effectiveness of bacteria acting as biological control agents in managing Fusarium head blight (FHB) will be conducted. Treatments will compare several bacterial strains used as biological control agents (BCAs) applied at various stages of crop development to an untreated control and a fungicide (Folicur) standard. A variety of BCAs will be cultured and applied from investigators in Illinois, Nebraska, and New York as part of a uniform BCA trial conducted in these other states as well. Strains of *Bacillus* isolated from South Dakota which have been studied in our lab for the last several years will also be used. Because it is difficult to guarantee FHB occurrence, the trial will be planted on sites with spring wheat residue and the site will be challenge inoculated with *Fusarium graminearum* conidia. Plots will be mist-irrigated to increase the likelihood of disease development. Foliar application of cells of the bacterial BCAs will be used in the field at flowering. Greenhouse trials using pots will also be conducted involving foliar spray application of selected BCAs at anthesis. Emphasis will be placed on greenhouse and field work combining one or more of our *Bacillus* isolates with the *Lysobacter* strain of Gary Yuen (U. of Nebraska) to see if a combination of the two BCAs will provide greater protection against FHB than either BCA alone. In addition, a commercial biocontrol product, AgraQuest's Serenade, will be used in greenhouse and field trials and compared to the efficacy of our South Dakota isolates as well as to Folicur. Combination formulations of Folicur and Serenade, and one or more BCAs will be studied to see if the combination of commercial products leads to greater control of FHB than use of each product in isolation.

Laboratory studies with pure cultures of the BCAs isolated from South Dakota wheat residue and foliage will examine the production of antibiotics (such as iturin) by these *Bacillus spp.* in a limited number of growth media. Effect of incubation time on antibiotic production and ability of the BCA inoculum to control FHB will be examined. This work is needed to optimize a growth medium and incubation time to optimize antibiotic production by BCAs whose antagonism of FHB relies at least in part on antibiotic production.

Bacterial BCAs may offer a more environmentally friendly control option for plant diseases than some chemical fungicides. Questions of health risks to the consumer are minimized with suitable BCAs. By using bacterial BCAs isolated from wheat in a local environment, compatibility of the agent with the crop or success in establishing the agent on the crop should be more reliable. Results of this study will provide guidance in how best to use the agents in a commercial wheat production system and how effective the BCAs may be in field situations.