

Report of the 2023 Uniform Regional Scab Nursery for Spring Wheat Parents

Report Prepared by:
Sarah Blecha and Andy Read
USDA-ARS
St. Paul, MN

The Uniform Regional Scab Nursery for Spring Wheat Parents (URSN) was grown for the 29th year in 2023. Four locations (St. Paul, MN, Crookston, MN, Brookings, SD, and Prosper, ND) reported results.

A total of 22 entries were included in the 2023 URSN, in addition to the resistant checks ND2710, BacUp, and Rollag, the susceptible checks Wheaton, Oslo, Norm, and N10, a Norm near-isoline containing *Fhb1*. The entries were contributed by four university wheat breeding programs.

The core set of traits evaluated at the nursery locations varied, but most included Fusarium head blight (FHB) incidence, FHB severity, and disease index. In addition, visual scabby kernel ratings (VSK/tombstone/FDK) were provided for locations. Additional agronomic trait data are presented in individual location summary tables for locations where they were measured. Adult plant and seedling stem rust reactions are also presented. Molecular marker genotypes for a set of FHB resistance QTL and other traits are provided for entries.

The data of genotype means for locations (Tables 2-5) will be available online on the Triticeae Toolbox under T3/Wheat.

This material is based upon work supported by the U.S. Department of Agriculture, in cooperation with the U.S. Wheat & Barley Scab Initiative. Any opinions, findings, conclusions, or recommendations expressed in this publication are those of the authors and do not necessarily reflect the view of the U.S. Department of Agriculture. USDA is an equal opportunity provider and employer.

St. Paul, MN
March 7, 2024

CONTENTS	PAGE
Cooperating Agencies, Stations and Personnel	1
Previous URSN Entries Released as New Cultivars	2-3
Figure 1. Geographic Locations of 2023 URSN	4
Table 1. List of Entries in the 2023 URSN	5
Tables 2-5. Nursery Data by Individual Location	6-9
Table 6. Summary of Means Across Locations	10
Table 7. Correlation Coefficients Between Traits by Location	11
Table 8. Correlation Coefficients Across Locations	12
Table 9. Seedling Stem Rust Reactions	13-14
Table 10. Molecular Marker Genotypes for Selected Genes and Traits	15-17
Table 11. Seedling and Field Stem Rust Reactions in St. Paul, Kenya and Ethiopia	18-23

Cooperators for the 2023 Uniform Regional Scab Nursery for Spring Wheat Parents

South Dakota State University (Brookings):

Karl Glover

University of Minnesota (St. Paul, Crookston):

Jim Anderson and Ruth Dill-Macky

North Dakota State University (Prosper, Langdon):

Andrew Green

USDA-ARS, Cereal Crops Research Unit (Fargo, ND):

Raj Nandety and Jason Fiedler

USDA-ARS, Cereal Disease Laboratory (St. Paul, MN):

Jim Kolmer, Yue Jin, and Matt Rouse

Previous URSN Entries Released as New Cultivars

Unlike the Hard Red Spring Wheat Uniform Regional Nursery (HRSWURN) Annual Report, the URSN Annual Report has not included records of experimental lines becoming released varieties. Including this information is valuable to the wheat community for tracking experimental lines that were tested in this nursery under FHB-inoculated conditions and their status of becoming a released variety. Some of these varieties were also tested in the HRSWURN as well, and information on their phenotype data can be found in that Annual Report posted to the GrainGenes website: <https://wheat.pw.usda.gov/GG3/>

The released cultivar names along with their experimental name, PI number, years in the URSN, and a brief description are included below. Entries that have been entered in the URSN since its start in 1995 are listed below in order of year it was released.

Briggs (SD3367, PI 632970) Released in 2002 by the South Dakota Agricultural Experiment Station. It has high and stable yield potential, above average test weight and protein content. It has good disease resistance to leaf rust and stem rust, and an intermediate resistance to FHB. **URSN 1998, HRSWURN 2000-2001**

Cromwell (MN00261-4, PI 653527) Released in 2007 by Thunder Seed, license required for seed production. Moderately strong straw, good test weight, and good end-use quality. Moderately resistant to FHB. **URSN 2006, HRSWURN 2004-2005**

Tom (MN01311-A-1, PI 656383) Released in 2008 by the University of Minnesota Agricultural Experiment Station due to its high yield, moderate resistance to FHB (type 2 resistance), and adult plant resistance to Ug99 race of stem rust. **URSN 2006, HRSWURN 2004-2005,**

Sabin (MN03358-4, PI 659083) Released in 2009 by the University of Minnesota Agricultural Experiment Station due to its high yield, moderate resistance to FHB due to the Fhb1 locus, and has the Lr34 gene for adult plant resistance to leaf rust. It heads relatively later than other cultivars in the region. **URSN 2008, HRSWURN 2006-2007**

Brick (SD3851, PI 657697) Released in 2009 by the South Dakota Agricultural Experiment Station due to its high resistance to FHB and high yield potential. It has early maturity and is moderately resistant to leaf rust. **URSN 2007, HRSWURN 2005-2006**

Select (SD3948, PI 659554) Released in 2010 by the South Dakota Agricultural Experiment Station due to its high resistance to FHB with high yield potential. It is also moderately resistant to leaf rust and has an early heading date. **URSN 2009, HRSWURN 2007-2008**

Rollag (MN05214-3, PI 665250) Released in 2011 by the University of Minnesota Agricultural Experiment Station due to its good resistance to FHB and has the Fhb1 locus. It also has a competitive yield, very good straw strength, and acceptable end-use quality. **URSN resistant check 2019-2023, HRSWURN 2009-2010**

Forefront (SD3997, PI 664483) Released in 2012 by the South Dakota Agricultural Experiment Station due to its competitive high yield, and protein content. It has resistance to FHB and leaf rust, along with an early heading date. **URSN 2009, HRSWURN 2009-2010**

Lanning (MT1316, PI 676978) Released in 2016 Montana Agricultural Experiment Station due to its high yield potential in dryland conditions and superior end-use quality. It is an awned, semidwarf hard red spring wheat. It is hollow-stemmed, making it susceptible to wheat stem sawfly. **URSN 2015-2016 and 2019, HRSWURN 2015-2016**

Lang-MN (MN10261-1, PI 687038) Released in 2017 by the University of Minnesota Agricultural Experiment Station. It has high yield and grain protein, good end-use quality, and disease resistance, especially to FHB. **URSN 2016, HRSWURN 2013-2014**

ND VitPro (ND825, PI 682660) Released in 2017 by the North Dakota Agricultural Experiment Station. It has improved straw strength, end-use quality, and good disease resistance, including FHB and stem rust. It has high vitreous kernel percentage and high grain protein. **URSN 2016, HRSWURN 2016**

Dagmar (MT1621, PI 690450) Released in 2019 by Montana Agricultural Experiment Station for adaptation to drought and wheat stem sawfly regions. It has a longer green leaf stage after heading, good grain protein, and good gluten strength. It has intermediate stem-solidness providing protection against wheat stem sawfly. **URSN 2018-2019, HRSWURN 2018-2019**

ND Frohberg (NDHRS16-13-97, PI 698310) Released in 2020 by the North Dakota Agricultural Experiment Station. It has improved straw strength, yield potential, and end-use quality. It has high resistance to stem rust, moderate resistance to leaf rust, bacterial leaf streak, and FHB. **URSN 2018, HRSWURN 2018-2019**

MT Sidney (MT1716, PI 699957) Released in 2021 by the Montana Agricultural Experiment Station. It is a high yielding, early-heading, awned, and semidwarf hard red spring wheat variety. Has moderate resistance to FHB. Hollow-stem makes it susceptible to wheat stem sawfly. **URSN 2019-2021, HRSWURN 2020-2021**

MT Dutton (MT1809) Released in 2023 by the Montana Agricultural Experiment Station. It has high yield in rainfed conditions, good protein content, and moderate aluminum tolerance. It is also moderately resistant to FHB and has resistance to foliar disease. **URSN 2020, HRSWURN 2021-2022**

ND Heron (NDHRS16-14-126, PI 699926) Released in 2022 by North Dakota State University. It has competitive yield potential with good protein, test weight, and resistance to FHB and bacterial blight. It has excellent end-use qualities. **URSN 2019-2020, HRSWURN 2019-2021**

Figure 1. Map of Uniform Regional Scab Nursery Locations, 2023




Table 1. Entries for the 2023 Uniform Regional Scab Nursery Parents.

Entry	Line	Pedigree	1st Year in URSN	Submitter	Organization
1	Bacup	CHECK			
2	ND2710	CHECK			
3	Rollag	CHECK			
4	Oslo	CHECK			
5	Wheaton	CHECK			
6	Norm	CHECK			
7	N10	CHECK (Norm Fhb1 NIL)			
8	MN18091-2	MN11116-3/MN12544-2	2023	J. Anderson	UMN
9	MN19114-3	MN13588-4/Lang-MN	2023	J. Anderson	UMN
10	MN19146-4	MN-Torgy/Lang-MN	2023	J. Anderson	UMN
11	MN19257-1	MN12345-3/MN-Washburn	2023	J. Anderson	UMN
12	MN20231-2	Shelly/MN-Torgy	2023	J. Anderson	UMN
13	SD5101	DRIVER/SD4708	2023	K. Glover	SDSU
14	SD5117	SY-VALDA/SD4748	2023	K. Glover	SDSU
15	SD5127	FOCUS/SD4710//SD4954	2023	K. Glover	SDSU
16	SD5132	SD4724/SD4719	2023	K. Glover	SDSU
17	SD5145	SD4752/SD4746	2023	K. Glover	SDSU
18	SD5159	SD4871/SD4539	2023	K. Glover	SDSU
19	NDHRS15-0037-C52	NDVITPRO/ROLLAG//GLENN ND820/LANG-	2023	A. Green	NDSU
20	NDHRS15-0006-B02	MN//DAPPS/3/GLENN/4/FALLER ND820/LANG-	2023	A. Green	NDSU
21	NDHRS15-0006-C29	MN//DAPPS/3/GLENN/4/FALLER	2023	A. Green	NDSU
22	NDHRS16RIL-0190-019	NDVITPRO/ND826	2023	A. Green	NDSU
23	NDHRS15-0405-C34	LANG-MN/ND829	2023	A. Green	NDSU
24	MT21214	DAGMAR/MT 1716	2023	J. Cook	MSU
25	MT21220	DAGMAR/MT 1716	2023	J. Cook	MSU
26	MT21222	DAGMAR/MT 1716	2023	J. Cook	MSU
27	MT21305	DAGMAR/NDHRS16-12-31	2023	J. Cook	MSU
28	MT21352	DAGMAR///MT1007//MO8/3-4	2023	J. Cook	MSU
29	MT21384	DAGMAR///MT1007//M0 09/3-4	2023	J. Cook	MSU

Table 2. 2023 Uniform Regional Scab Nursery for Spring Wheat Parents, St. Paul, MN.

Entry	Line	Incidence %	Severity %	Disease Index	VSK %	DON ppm	Heading d from 6-1	micro TWT ¹ g
1	Bacup	83.3	13.0	11.1	9.0	2.5	20.0	11.4
2	ND2710	69.0	8.5	6.1	6.0	1.2	21.0	11.7
3	Rollag	83.3	16.0	14.1	9.0	2.6	21.0	11.2
4	Oslo	95.2	40.5	39.4	13.5	3.2	21.0	10.3
5	Wheaton	100.0	80.8	80.8	77.5	17.7	23.0	9.1
6	Norm	100.0	66.7	66.7	62.5	12.2	24.3	10.0
7	N10	100.0	39.7	39.7	38.5	8.3	25.7	10.4
8	MN18091-2	90.4	26.3	24.1	12.5	4.7	27.0	11.5
9	MN19114-3	100.0	20.2	20.2	12.0	2.0	24.3	11.5
10	MN19146-4	78.5	10.5	8.5	8.0	1.8	23.0	11.7
11	MN19257-1	92.8	16.7	15.6	6.0	1.3	27.0	11.7
12	MN20231-2	52.4	8.6	4.3	11.5	2.3	23.0	11.7
13	SD5101	85.7	17.1	15.5	9.0	2.7	22.0	11.6
14	SD5117	80.9	17.0	14.2	10.0	1.2	20.0	11.4
15	SD5127	83.3	25.0	20.4	16.0	3.6	21.0	10.8
16	SD5132	97.6	31.4	30.5	16.0	2.2	27.0	11.4
17	SD5145	97.6	16.5	16.1	6.0	0.8	21.0	11.7
18	SD5159	90.4	30.4	28.3	16.0	2.8	23.0	11.4
19	NDHRS15-0037-C52	95.2	24.6	23.4	12.5	3.7	23.0	11.9
20	NDHRS15-0006-B02	83.3	14.1	12.7	7.0	2.6	20.0	11.4
21	NDHRS15-0006-C29	88.1	15.0	13.2	7.0	2.5	20.0	11.7
22	NDHRS16RIL-0190-019	78.5	21.8	17.0	12.5	4.3	20.0	11.5
23	NDHRS15-0405-C34	95.2	23.2	22.4	15.0	3.6	21.0	11.2
24	MT21214	97.6	32.0	31.8	20.0	5.5	24.3	11.0
25	MT21220	100.0	34.9	34.9	12.5	3.1	25.7	11.5
26	MT21222	95.2	24.9	23.7	16.0	4.7	24.3	11.3
27	MT21305	95.2	15.2	14.3	7.0	3.4	20.0	11.6
28	MT21352	100.0	29.9	29.9	30.0	9.3	20.0	10.8
29	MT21384	95.2	30.3	29.6	16.0	7.8	20.0	11.0
*	Alsen	97.6	21.8	21.5	11.0	2.7	22.0	11.4
*	Roblin	97.6	61.3	60.3	32.5	5.1	20.0	10.9
*	MN00269	100.0	55.3	55.3	32.5	3.0	27.0	10.1
Mean		90.6	27.8	26.4	17.8	4.2	22.6	11.2
LSD		24.2	22.4	22.8	13.8	-	2.1	0.5
CV		13.0	39.3	42.1	37.8	-	5.8	2.2

¹ Weight of the VSK sample that fits in a 15.7 mL copper vessel measuring 20 mm in diameter and 50 mm in height

* Extra entries

Table 3. 2023 Uniform Regional Scab Nursery for Spring Wheat Parents, Crookston, MN.

Entry	Line	Incidence ¹	Severity ¹	Disease ¹	VSK	DON	Heading	micro TWT ²
		%	%	Index	%	ppm	d from 6-1	g
1	Bacup	–	–	–	10.0	6.6	32.0	11.5
2	ND2710	–	–	–	8.0	6.7	36.3	11.8
3	Rollag	–	–	–	12.5	7.7	36.3	11.1
4	Oslo	–	–	–	65.0	11.5	34.0	9.7
5	Wheaton	–	–	–	70.0	26.8	38.3	9.1
6	Norm	–	–	–	47.5	21.9	38.3	10.1
7	N10	–	–	–	26.0	16.3	37.0	10.6
8	MN18091-2	–	–	–	13.5	13.1	41.3	11.4
9	MN19114-3	–	–	–	14.5	17.0	42.3	11.1
10	MN19146-4	–	–	–	7.0	8.3	37.7	11.9
11	MN19257-1	–	–	–	8.0	6.8	43.0	11.4
12	MN20231-2	–	–	–	12.5	7.9	33.0	11.2
13	SD5101	–	–	–	13.5	11.5	37.3	11.4
14	SD5117	–	–	–	13.5	5.0	35.0	11.3
15	SD5127	–	–	–	14.5	10.2	30.7	11.5
16	SD5132	–	–	–	9.0	10.4	43.0	11.4
17	SD5145	–	–	–	10.0	6.8	38.3	11.6
18	SD5159	–	–	–	9.0	5.4	38.3	11.7
19	NDHRS15-0037-C52	–	–	–	8.0	10.3	37.0	12.0
20	NDHRS15-0006-B02	–	–	–	9.0	11.7	33.3	11.4
21	NDHRS15-0006-C29	–	–	–	16.0	11.4	35.0	11.4
22	NDHRS16RIL-0190-019	–	–	–	9.0	10.1	35.0	11.7
23	NDHRS15-0405-C34	–	–	–	11.0	9.3	34.3	11.3
24	MT21214	–	–	–	35.0	26.6	39.7	10.5
25	MT21220	–	–	–	17.0	17.3	40.3	11.3
26	MT21222	–	–	–	12.0	9.1	41.3	11.3
27	MT21305	–	–	–	13.5	10.1	34.0	11.1
28	MT21352	–	–	–	37.5	22.7	35.3	10.6
29	MT21384	–	–	–	20.0	19.3	36.0	10.3
*	Alsen	–	–	–	8.0	8.3	38.7	11.8
*	Roblin	–	–	–	77.5	12.2	32.0	9.7
*	MN00269	–	–	–	40.0	25.2	45.3	9.5
Mean		–	–	–	21.2	12.6	37.2	11.0
LSD		–	–	–	14.4	-	1.4	0.6
CV		–	–	–	33.2	-	2.4	2.6

¹ Not collected- symptoms were slow to develop and by the time they worth scoring, later heading lines had senesced too far

² Weight of the VSK sample that fits in a 15.7 mL copper vessel measuring 20 mm in diameter and 50 mm in height

* Extra entries

Table 4. 2023 Uniform Regional Scab Nursery for Spring Wheat Parents, Brookings, SD.

Entry	Line	Incidence %	Severity %	Disease %	Tombstone %
1	Bacup	80.0	16.3	13.0	22.5
2	ND2710	76.4	9.8	7.5	7.5
3	Rollag	95.0	18.3	17.5	27.5
4	Oslo	95.0	18.5	17.7	36.7
5	Wheaton	100.0	25.3	25.3	58.3
6	Norm	87.5	15.3	13.6	20.0
7	N10	92.5	17.0	16.1	30.0
8	MN18091-2	93.3	16.2	15.2	12.3
9	MN19114-3	95.0	15.5	14.8	13.5
10	MN19146-4	80.0	11.5	9.5	16.7
11	MN19257-1	87.5	13.5	12.0	5.0
12	MN20231-2	95.0	16.3	15.5	9.7
13	SD5101	90.0	16.5	15.0	15.0
14	SD5117	96.7	18.7	18.1	13.3
15	SD5127	90.0	14.8	13.7	22.5
16	SD5132	92.2	14.4	13.3	9.0
17	SD5145	93.3	18.0	16.8	15.0
18	SD5159	91.7	17.0	16.2	13.0
19	NDHRS15-0037-C52	88.3	16.0	14.4	8.3
20	NDHRS15-0006-B02	74.5	11.4	8.5	15.0
21	NDHRS15-0006-C29	97.5	18.8	18.3	10.0
22	NDHRS16RIL-0190-019	90.0	15.0	13.6	16.7
23	NDHRS15-0405-C34	90.0	13.8	12.5	17.5
24	MT21214	90.0	17.0	15.5	26.7
25	MT21220	91.7	14.0	13.0	25.0
26	MT21222	96.7	17.3	16.9	25.0
27	MT21305	91.7	18.3	17.1	23.3
28	MT21352	90.6	22.8	20.6	27.5
29	MT21384	100.0	23.5	23.5	25.0
Mean		91	16.6	15.4	19.7
LSD 0.05		NS	6.10	7.6	10.3
CV		8.96	20	26.8	28.4

Field was very dry after planting which decreased germination. After eventual rain, weeds overtook some rows. Thirteen rows were entirely lost to weeds and ratings were collected from eight rows on fewer than 20 heads

Table 5. 2023 Uniform Regional Scab Nursery for Spring Wheat Parents, Prosper, ND.

Entry	Line	VIBE FDK ¹	FHB	Flowering date	DON	Notes
		%	9-Jan	DAP	ppm	
1	Bacup	22.3	6.2	203.5	8.2	
2	ND2710	17.5	4.6	202.2	6.3	
3	Rollag	17.6	6.0	202.0	6.5	
4	Oslo	12.4	5.9	202.1	4.5	
5	Wheaton	16.3	5.1	202.5	9.2	
6	Norm	46.6	6.9	202.6	28.1	
7	N10	41.0	6.2	202.0	23.6	
8	MN18091-2 *	-	-	-	-	
9	MN19114-3	15.7	4.9	202.8	15.7	
10	MN19146-4	15.7	4.6	202.5	5.3	
11	MN19257-1	9.9	2.9	203.4	14.9	
12	MN20231-2	16.6	4.5	201.7	7.6	
13	SD5101	17.9	5.2	203.1	6.7	
14	SD5117	12.8	5.0	201.8	2.1	
15	SD5127	11.0	7.8	202.5	4.6	
16	SD5132	25.8	4.0	203.4	18.9	
17	SD5145	15.6	5.1	203.7	9.6	
18	SD5159	9.7	4.4	204.0	7.1	
19	NDHRS15-0037-C52	11.5	5.7	203.8	8.8	
20	NDHRS15-0006-B02	16.0	4.3	203.0	5.3	
21	NDHRS15-0006-C29	11.5	5.3	202.9	5.3	
22	NDHRS16RIL-0190-019	19.3	6.7	202.8	7.3	
23	NDHRS15-0405-C34	14.5	6.4	203.0	7.0	
24	MT21214	23.4	6.1	202.0	10.0	
25	MT21220	25.4	5.6	202.5	12.5	
26	MT21222	22.8	3.7	203.0	21.9	
27	MT21305	16.7	6.2	203.4	8.4	
28	MT21352	15.5	7.7	202.2	9.3	
29	MT21384	14.2	5.3	201.6	10.5	
Mean		18.5	5.5	202.7	10.7	
CV		18.3	11.6	0.2	26.8	
LSD 0.05		6.6	1.3	0.9	6.9	
**	AKFASTRO	55.4	6.7	203.6	58.4	Sus Late
**	ALSEN	17.9	4.9	201.2	8.1	Res Med
**	GLENN	10.2	3.9	201.1	3.6	Res Early
**	LINKERT	28.8	6.8	203.1	7.3	Ms Late
**	ND828	7.4	4.9	202.7	9.7	R Late
**	NDFROHBERG	13.8	6.7	203.0	7.6	MR Med
**	NDHERON	9.9	6.5	203.0	1.2	MR Early
**	NDVITPRO	9.4	6.2	202.9	6.2	MR Med
**	WBMAYVILLE	18.7	9.0	203.6	3.2	Sus Med
**	2398	36.0	5.9	202.6	32.5	Sus Early

* Did not germinate.

** Local checks

¹ Fusarium Damaged Kernels calculated from RGB image of plot grain sample using algorithm from VIBE seed analyzer.

Table 6. 2023 Uniform Regional Scab Nursery for Spring Wheat Parents - Summary of Means.

Line	Incidence		Severity		Disease		VSK ^{1, 2}		DON ²	
	%	Rank	%	Rank	Index	Rank	%	Rank	ppm	Rank
	No. of Locations > 2		2		2		4		3	
Bacup	81.7	5	14.6	5	12.0	5	15.9	18	5.8	7
ND2710	72.7	1	9.2	1	6.8	1	9.8	2	4.7	2
Rollag	89.2	10	17.2	10	15.8	11	16.7	20	5.6	5
Oslo	95.1	21	29.5	27	28.6	27	31.9	26	6.4	10
Wheaton	100.0	29	53.0	29	53.0	29	55.5	29	17.9	28
Norm	93.7	18	41.0	28	40.2	28	44.1	28	20.7	29
N10	96.2	26	28.4	26	27.9	26	33.9	27	16.1	27
MN18091-2	91.9	14	21.3	19	19.7	18	12.8	11	8.9	19
MN19114-3	97.5	27	17.9	13	17.5	16	13.9	13	11.6	22
MN19146-4	79.3	4	11.0	2	9.0	2	11.9	7	5.1	4
MN19257-1	90.2	11	15.1	6	13.8	6	7.2	1	7.7	18
MN20231-2	73.7	2	12.5	3	9.9	3	12.6	10	5.9	8
SD5101	87.8	8	16.8	8	15.3	7	13.9	12	7.0	14
SD5117	88.8	9	17.8	12	16.2	12	12.4	9	2.8	1
SD5127	86.7	7	19.9	16	17.1	14	16.0	19	6.1	9
SD5132	94.9	20	22.9	20	21.9	20	15.0	16	10.5	20
SD5145	95.4	23	17.3	11	16.5	13	11.6	5	5.7	6
SD5159	91.1	12	23.7	21	22.3	21	11.9	8	5.1	3
NDHRS15-0037-C52	91.8	13	20.3	17	18.9	17	10.1	3	7.6	17
NDHRS15-0006-B02	78.9	3	12.8	4	10.6	4	11.7	6	6.5	12
NDHRS15-0006-C29	92.8	16	16.9	9	15.8	10	11.1	4	6.4	11
NDHRS16RIL-0190-019	84.3	6	18.4	14	15.3	8	14.4	14	7.2	15
NDHRS15-0405-C34	92.6	15	18.5	15	17.5	15	14.5	15	6.6	13
MT21214	93.8	19	24.5	23	23.7	22	26.3	24	14.0	26
MT21220	95.8	24	24.4	22	23.9	23	20.0	23	11.0	21
MT21222	96.0	25	21.1	18	20.3	19	18.9	22	11.9	23
MT21305	93.5	17	16.7	7	15.7	9	15.1	17	7.3	16
MT21352	95.3	22	26.4	24	25.3	24	27.6	25	13.8	25
MT21384	97.6	28	26.9	25	26.6	25	18.8	21	12.5	24

¹ FDK and Tombstone ratings are included, these terminologies are interchangeable

² One fewer location was used for MN18091-2 because of poor germination

Table 7. Correlation Coefficients Between Traits, by Location.

Correlation Between	St. Paul	Brookings	Crookston	Prosper
Incidence & Severity	0.571	0.743		
Incidence & Disease Index	0.614	0.834		
Incidence & Tombstone/VSK/FDK	0.395	0.354		
Incidence & DON	0.422			
Severity & Disease Index	0.997	0.986		
Severity & Tombstone/VSK/FDK	0.924	0.629		
Severity & DON	0.878			
Disease Index & Tombstone/VSK/FDK	0.919	0.611		
Disease Index & DON	0.874			
Tombstone/VSK/FDK & DON	0.944		0.699	0.787

Table 8. Correlation coefficients among traits, using means across locations

	Incidence	Severity	Disease Index
Severity	0.497		
Disease Index	0.551	0.997	
Tombstone/VSK/FDK	0.386	0.727	0.733

Calculated using 2 locations: St. Paul and Brookings

**Table 9. 2023 Uniform Regional Scab Nursery for Spring Wheat Parents, St. Paul, MN.
Seedling stem rust reactions (Y. Jin, USDA-ARS).**

Entry	Line	Race					Field stem rust response St. Paul
		QFCSC 06ND76C	QTHJC 75ND717C	RKRQC 99KS76A-1	TPMKC 74MN1409	TTTTF 01MN84A-1-2	
1	Bacup	0;	2	2-	0;	3	10R
2	ND2710	0	2-	;2-	;2-	1;	20R-MR
3	Rollag	0;	2-	2	2-2	1;	10R
4	Oslo	;	2	0;	2-	3	30MR
5	Wheaton	0	2-	;2-	2-	1;	5R
6	Norm	0	2-	;2-	2-	0;1	0
7	N10	0	2-	;2-	2-	1;	5R
8	MN18091-2	0;	2-	2-	2-	1	10R
9	MN19114-3	0;	2-	2	0;	1;	0
10	MN19146-4	0;	2-	2	0;	1;	5R
11	MN19257-1	0;	2-	2	2-	11+	0
12	MN20231-2	0	2-	2	0;	3-	5R
13	SD5101	;	2-C	2-	;	1+	40MR
14	SD5117	;	2-C	2-	;	11+	20MR
15	SD5127	;	2-C	2	2	3+	30MR
16	SD5132	0;	2-C	2/3	3	1;	0/30MS
17	SD5145	;	2-C	2+2	2	11+/2-	5R
18	SD5159	;	2-	2	;	;	10R
19	NDHRS15-0037-C52	;	2-	2	2-	1;	5R
20	NDHRS15-0006-B02	;	2-	2	2-	1;	5R
21	NDHRS15-0006-C29	;	2-	2	2	1;	5R
22	NDHRS16RIL-0190-019	;	2-	2	2-	1;	0
23	NDHRS15-0405-C34	;	2-	2	2-	1;	5R
24	MT 21214	;	2	22+	;2-/2	4	10R-MR
25	MT 21220	;	12-	2	;2-/2	3+	0
26	MT 21222	;	2-2	2	;2-/2	3+	5R
27	MT 21305	;2-	2-	2	2-	3+	20R-MR
28	MT 21352	;	2-	2	2	4	20R-MR
29	MT 21384	;2-	2-	2	2	3	40MR
*	Line E	33+	3	3+	4	4	90S
*	Line E	33+	3+	4	4	3+	90S
*	LMPG-6	2+2	2	2+3	3	2+3	90S
*	NA101/MqSr7a	1+1	23	1;	3+	1;	70MS-S

* checks

Explanatory notes on next page

Table 9 continued, Explanatory notes.

A. Races used in seedling evaluations:

Race	Origin	Virulence on differential genes
QFCSC	USA	5 8a 9a 9d 9g 10 17 21 McN
QTHJC	USA	5 6 8a 9b 9d 9g 10 11 17 21 McN
RKRQC	USA	5 6 7b 8a 9a 9b 9d 9g 17 21 36 McN
TPMKC	USA	5 7b 8a 9d 9e 9g 10 11 17 21 36 Tmp McN
TTTTF	USA	5 6 7b 8a 9a 9b 9d 9e 9g 10 11 17 21 30 36 38 Tmp McN

* **Red font** represents unique and/or significant virulence or combination of virulences to resistance genes that are important in spring wheat

B. Seedling rating scale:

0 to 4 infection type scale of Stakmen et al., 3 or 4 are considered susceptible

"H" denotes heterogeneous, the predominant type given first.

"LIF" denotes low infection frequency, or fewer number of pustules.

"C" stands for excessive chlorosis

"N" stands for excessive necrosis

"Sr2M" referred to seedling chlorosis, similar to Sr2 expression in seedling under certain environments

C. Field stem rust nursery:

Entries were planted in 1-m row plots

Nursery was inoculated with a bulk of races QFCSC, QTHJC, RCRQC, RTQQC, and TPMKC

Stem rust terminal severity (%) and infection responses (R, MR, MS, S or combination thereof) were rated when entries were at the soft dough stage

Table 10. Markers Associated With Selected Traits/Genes (R. Nandety and J. Fiedler, USDA-ARS).

Trait	Marker
StemRust 3B	Sr2
StemRust 6A	Sr6
StemRust 3B	Sr12
Stem Rust 7D	Sr25
LeafRust 2B	Lr13
LeafRust 2B	Lr16
LeafRust 1D	Lr21
LeafRust 2B	Lr23
LeafRust 7D	Lr34
YellowRust 2B	Yr7D
TanSpot 5B	Tsn1
Fhb 3B	Fhb1
Fhb 3B	Fhb1-TaHRC
Fhb 5A	barc180
Fhb 5A	barc186
Fhb 6B	gwm644
GrainProt. 6B	GPC
Glutenins 1D	GluD1
Glutenin 1A	um19
Dwarfing 4B	RhtB1
Dwarfing 4D	RhtD1
Dwarfing 6A	Rht24
Photoper. 2B	PpdB1
Photoper. 2D	PpdD1

Entry Line	Marker	StemRust 3B	StemRust 6A	StemRust 3B	Stem Rust 7D	LeafRust 2B	LeafRust 2B	LeafRust 1D	LeafRust 2B	LeafRust 7D	YellowRust 2B	TanSpot 5B	Fhb 3B	Fhb 3B	Fhb 5A	Fhb 5A	Fhb 6B	GrainProt. 6B	Glutenins 1D	Glutenin 1A	Dwarfing 4B	Dwarfing 4D	Dwarfing 6A	Photoper. 2B	Photoper. 2D
1	Bacup	S	S	R	R	R	S	S	S	R	S	S	S	S	R	R	S	N	G	1	Wt	Wt	D	I	I
2	ND2710	S	S	R	S	R	S	S	R	R	R	S	R	R	R	R	S	N	G	1	Wt	Wt	Wt	S	S
3	Rollag	S	S	R	R	R	R	S	R	R	S	R	R	R	S	R	S	N	G	2	Wt	D	D	S	S
4	Oslo	S	R	R	S	R	S	S	S	R	R	S	S	S	S	S	S	N	P	1	D	Wt	D	S	I
5	Wheaton	S	S	R	S	R	S	S	R	R	R	R	S	S	S	S	S	N	G	2	Wt	D	Wt	S	I
6	Norm	S	S	R	S	R	R	S	R	R	R	R	S	S	S	S	S	N	G	2	Wt	D	Wt	S	S
7	N10	S	S	R	S	R	S	S	R	R	R	R	R	R	S	S	S	N	P	1	Wt	D	Wt	S	S
8	MN18091-2	-	S	R	S	S	S	S	R	R	S	R	R	R	S	S	S	N	G	2	D	Wt	Wt	S	S
9	MN19114-3	-	S	R	R	R	S	S	R	R	R	S	R	R	S	S	S	N	G	2	Wt	D	D	S	S
10	MN19146-4	-	S	R	R	R	S	S	R	R	R	S	R	R	S	S	S	N	G	1	Wt	Wt	D	S	I
11	MN19257-1	S	S	R	S	R	S	S	S	R	S	R	R	R	R	R	S	N	G	2	D	Wt	D	S	S
12	MN20231-2	-	S	R	R	S	S	R	R	S	R	R	R	R	S	S	S	N	G	2	Wt	Wt	D	S	S
13	SD5101	-	S	S	S	S	R	S	R	S	R	R	S	S	S	S	S	N	G	2	Wt	Wt	D	S	S
14	SD5117	S	S	S	S	S	R	S	S	R	R	S	S	S	S	S	S	N	G	2	Wt	Wt	Wt	S	S
15	SD5127	-	S	R	R	R	R	R	R	S	R	R	R	R	S	S	S	N	G	2	D	Wt	Wt	S	I
16	SD5132	S	S	R	S	S	R	R	S	R	R	S	S	S	S	S	S	N	G	2	Wt	Wt	D	S	S
17	SD5145	S	S	R	S	R	R	S	R	S	R	S	S	S	S	S	S	N	G	2	Wt	Wt	D	S	S
18	SD5159	S	S	S	S	S	S	R	S	R	R	R	R	R	S	S	S	N	G	2	Wt	Wt	D	S	I
19	NDHRS15-0037-C52	-	S	R	S	R	R	R	S	R	S	S	S	S	S	S	S	N	G	2	D	Wt	Wt	S	S
20	NDHRS15-0006-B02	-	S	R	R	S	S	S	R	S	S	S	S	S	R	R	S	N	G	2	D	Wt	Wt	S	S
21	NDHRS15-0006-C29	-	S	R	S	S	S	S	R	S	S	R	R	R	R	R	S	N	G	2	D	Wt	Wt	S	S
22	NDHRS16RIL-0190-019	-	S	R	S	R	R	R	S	R	S	S	S	S	S	S	S	N	G	2	D	Wt	Wt	S	S
23	NDHRS15-0405-C34	-	S	R	S	R	S	R	S	R	R	S	S	S	S	S	S	N	G	2	D	Wt	D	S	I
24	MT21214	-	S	R	S	S	S	S	S	R	R	S	S	S	S	S	S	N	G	2	D	Wt	Wt	S	S
25	MT21220	-	S	R	S	R	R	S	R	S	R	R	S	S	S	S	S	N	G	2	U	Wt	Wt	S	S
26	MT21222	-	S	R	S	R	S	S	R	S	R	R	S	S	S	S	S	N	G	2	D	Wt	Wt	S	S
27	MT21305	-	S	R	S	R	R	S	R	S	R	R	S	S	R	R	S	N	G	2	D	Wt	Wt	S	S
28	MT21352	S	S	R	S	S	S	S	S	R	R	S	S	S	S	S	S	N	G	1	D	Wt	Wt	S	S
29	MT21384	-	S	R	S	S	R	S	R	S	R	S	S	S	S	S	S	N	G	2	D	Wt	Wt	S	S

Information about markers on next page

Allele Code	Description
R = Resistant (Hope allele)	S = Susceptible
R = Resistant (Harvest allele)	S = Susceptible
R = Resistant	S = Susceptible
R = Resistant (200 bp present)	S = Susceptible (no 200 bp)
R = Ne2/Lr13_ha, Lr13 resistant and hybrid necrosis (Ne2 allele)	S = Ne2/Lr13_ha, Lr13 susceptible, hybrid necrosis only but still has Ne2
R = Resistant	S = Susceptible
R = Resistant	S = Susceptible
R = Resistant	S = Susceptible
R = Resistant	S = Susceptible
R = Resistant (Thatcher allele)	S = Susceptible
R = Resistant	S = Susceptible
R = Resistant	S = Susceptible
R = Resistant	S = Susceptible
R = Resistant	S = Susceptible
R = Resistant	S = Susceptible
R = Resistant	S = Susceptible
R = Resistant	S = Susceptible
R = Resistant	S = Susceptible
R = Resistant (161 bp present)	S = Susceptible (no 161 bp)
I = Increased	N = Normal
G = Good (5+10)	P = Poor (2+12)
I = 359bp = Ax1 or Ax-null	2 = 341bp = Ax2
D = Dwarfing = Rht-B1b	Wt = Wild Type = Rht-B1a
D = Dwarfing = Rht-D1b	Wt = Wild Type = Rht-D1a
D = Dwarfing	Wt = Wild Type
I = Insensitive	S = Sensitive
I = Insensitive	S = Sensitive

U = No Call or Unknown = Indeterminant designation

Het = Heterozygous call

Table 10 continued, Marker information

Name	Chromosome	Alternate Name	Comment	Manuscript
Sr2	3B		Null allele	https://doi.org/10.1007/s00122-010-1482-7
Sr8	6A	kwh53		https://doi.org/10.1094/PHYTO-05-16-0186-R
Sr12	3B	NBLRR3		https://doi.org/10.1371/journal.pone.0157029
Sr25	7D		SSR	Chao, unpublished https://www.cell.com/molecular-plant/pdf/S1674-2052(21)00171-4.pdf
Lr13	2B			https://doi.org/10.1186/s12870-017-0993-7
Lr16	2B	kwm849		https://doi.org/10.1007/s11032-012-9773-0
Lr21	1D			https://doi.org/10.1007/s11032-017-0628-6
Lr23	2B	sunKASP_16 FJ436983-		https://doi.org/10.1126/science.1166453
Lr34	7D	T67957A		https://doi.org/10.1038/s41477-018-0236-4
Yr7	2B	Yr7D		
Tsn	5B	Tsn1-1Ka	SNP flanking deletion	Faris Lab unpublished
Fhb1	3B	FM227		https://doi.org/10.1007/s00122-016-2727-x
TaHRC	3B			https://doi.org/10.1007/s00122-018-3159-6
barc180	5A	GENE-3371_56	equivalent to SSR	https://doi.org/10.1007/s00122-011-1573-0
barc186	5A	IWA6412	equivalent to SSR	Chao, unpublished
gwm644	6B		SSR	https://doi.org/10.1093/genetics/149.4.2007
GPC	6B	GPC-B1_DUP		https://doi.org/10.1111/j.1469-8137.2005.01627.x
GluD1	1D			https://doi.org/10.1270/jsbbs.57.243
umn19	1A		SSR	https://doi.org/10.1007/s00122-008-0886-0
RhtB1	4B			https://doi.org/10.1007/s00122-002-1048-4
RhtD1	4D			https://doi.org/10.1007/s00122-002-1048-4
Rht24	6A	Rht24-TaAP2.1		Anderson lab
PpdB1	2B			https://doi.org/10.1371/journal.pone.0079459
PpdD1	2D			https://doi.org/10.1007/s11032-012-9765-0

